

WL IRON CHELATE

ANALYSIS
IRON (Fe)
SULPHUR (S)

W/W
5%
4%

Derived from: Ferrous Sulphate and Phenolic Acid Chelate

PRODUCT CONTAINS 76% W/W WATER
TOTAL SOLIDS 24% W/W

GENERAL INFORMATION

- WL IRON Chelate is a foliar or soil applied micronutrient and is nonphytotoxic when used as directed. It is completely available and absorbed by the leaf surface or the root system because of the natural complexing properties of its phenolic acid chelating complex.
- It is an excellent acidification agent for lowering high pH in spray tank water to decrease the hydrolysis of most pesticides.
- It is an effective dispersing agent when applied with other foliar spray chemicals.

MIXING INSTRUCTIONS

- WL IRON Chelate will disperse in water with little agitation. Many pesticides can be added and applied while spraying. Refer to pesticide label.

When foliar spraying WL IRON Chelate through conventional sprayers, use a minimum of 200 litres of water per hectare. When spraying with low volume equipment, 50 litres of water per hectare is usually sufficient. If less water is used, slight burning of the foliage may occur. A maximum of 4 litres per hectare per application is recommended. Aerial applications should not exceed 1 litre per 4 litres of water. For tree crops and dilute sprays refer to table opposite. WL IRON Chelate is manufactured to rigid controls at the highest possible concentration for effective usage. Any residue in this pail is water soluble. Rinse the pails with water and add solution to spray tank.

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

CAUTION: Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

BMP1016

Stoller's



IRON Chelate



Stoller

STOLLER AUSTRALIA PTY. LTD.

A.C.N. 065 320 747

1 Creswell Road
Largs Bay SA 5016
Australia

Phone 08 8169 0900
Fax 08 8169 0909
www.stoller.com.au

CONTENTS 10 20 200 800 1000 LITRES

MADE IN AUSTRALIA

CAUTION

Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Use gloves and goggles.

DIRECTIONS FOR USE

GENERAL APPLICATION RATES WHEN LEAF & SOIL TEST ARE NOT AVAILABLE

	SOIL APPLICATION	
Maintenance application		2 litres per hectare
Moderate deficiency		4-9 litres per hectare
Heavy micronutrient deficiency		9-18 litres per hectare

For best results spray when crop is in an active growing state, after irrigation or natural rain fall. Spray early in the morning or late in the afternoon. Midday sprays may not be effective because of excessive moisture evaporation. The addition of .5% (total solution) of nitrogen solution, ammonium sulphate or L.B. Urea may aid leaf absorption. Do not spray in hot weather conditions.

FOLIAR SPRAY

CROP	RATE	COMMENTS
Vegetables	1 - 2 litres per hectare each spray	Incorporate WL IRON Chelate into the normal spray program. Frequent applications may be required depending on severity of deficiency.
Trees, Fruits Nuts and Grapevines	2 - 4 litres per hectare	If tree crops are sprayed several times a year, the amount can be split into more numerous applications. Split applications are considered more beneficial than single applications.
Field Crops	2 - 4 litres per hectare	Apply recommended rates when plants are young. This is the time when most severe deficiencies occur.

TRICKLE IRRIGATION

Apply to trees and vines through irrigation at 4-6 litres per hectare per application.

LIQUID FERTILIZER

Mix WL IRON Chelate with liquid fertilizer agitating to thoroughly disperse the trace elements. This can be achieved by adding the product to the mixing batch tank or inducting through a pump. When mixing in batch tank, add product to water or nitrogen solution before adding phosphate material.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No: