

ACTION 10

ANALYSIS

CALCIUM (Ca) as chloride

Formulated with Stoller's proprietary mix of Stoller's Co-Factors.

%W/V

10 %

GENERAL INFORMATION

- ACTION 10 is a secondary nutrient combined with natural growth activators. ACTION 10, in conjunction with a good nutritional program, provides an extra boost to seedling establishment and root development.
- ACTION 10 provides an important boost to any plant's tolerance to adverse conditions and stresses. Action 10 will enhance crop growth when temperatures prevail below 21°C or above 31°C and plant growth is suspended.
- ACTION 10 is completely available and absorbed by the leaf surface or the root system.
- ACTION 10 is designed for use in foliar, seed row, seed and transplant water application and is non-phytotoxic when used as directed.

MIXING INSTRUCTIONS

ACTION 10 will disperse in water with little agitation. Many other agricultural chemicals can be sprayed simultaneously.

CAUTION: Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

WARNING: Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Use gloves and goggles.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

BMP0517



Stoller

ACTION 10

STOLLER AUSTRALIA PTY. LTD.

1 Creswell Road
Largs Bay SA 5016

Phone: 08 8169 0900
Fax: 08 8169 0909
www.stoller.com.au

Made in Australia

CONTENTS 5 10 20 200 1000 LITRES

DIRECTIONS FOR USE

CROP	RATE	COMMENTS
Cotton	1.0 litre per hectare	Apply in furrow at planting directly onto the seed where Black Root Rot is a problem
	1.0 litre per hectare	Apply in furrow at planting for improved germination and establishment
	500 mls - 1 litre per hectare	Apply as a foliar spray if growing/weather conditions persist that are conducive to Black Root Rot or slow growth
<i>This can be applied as a band spray as soon as the crop target is big enough. The 500 mls per hectare rate should be used only when 1.0 litre per hectare has already been used in furrow at planting. At all other times use the 1.0 litre per hectare rate.</i>		
Sugarcane	500 mls - 1 litre per hectare	Apply in furrow at planting to encourage vigour. Apply as a foliar spray 0-45 days post harvest.
Wheat, Barley, Corn, Green Beans, French Beans	500 mls per hectare	Apply in furrow at planting or as a foliar spray as soon as the plant is a big enough target.
	1.0 litre per hectare	Apply as a foliar spray for improved growth after planting or when adverse weather conditions persist. Apply in furrow at planting directly onto the seed where Red Root Rot is a problem.
Faba Beans, Chick Peas, Soybeans	500 mls per hectare	Apply as a foliar spray from first flower bud to first full flowers open

Potatoes and All other crops Consult your Agronomist/Consultant or your Stoller Australia Distributor

SEED TREATMENT (General)

To improve seedling establishment and early vigour, apply ACTION 10 to the seed at 500 ml to 1.0 litres per 100kgs of seed (grain). Apply ACTION 10 to potato seed at 2 litres per tonne of seed as a dip, or 5L per hectare in furrow directly over the seed. ACTION 10 is water based and should not be mixed with oil based seed treatments. Allow seed to dry before applying oil based treatments. Do not use treated seed for food, feed or oil purpose.

IN FURROW APPLICATION (General)

ACTION 10 can be used in-furrow at the time of seeding or planting and should be applied at a rate of 500 ml to 1.0 litres per hectare.

FOLIAR APPLICATION (General)

For continued healthy root growth and plant integrity apply ACTION 10 at regular intervals at a rate of 500 ml to 1.0 litres per hectare. Repeat applications every 7 – 14 days. Dilution rates should not exceed 2% (i.e. 1 litre of ACTION 10 should be mixed with a minimum of 50 litres of water). Mix 10 ml of ACTION 10 per litre of water for hand spraying.