

## CANOPY MASTER 10

### ANALYSIS

	W/W
Nitrogen (as urea)	5.0%
Magnesium (as magnesium sulphate)	3.0%
Zinc (as zinc sulphate)	2.0%
Manganese (as manganese sulphate)	2.0%
Iron (as iron sulphate)	1.0%
Copper (as copper sulphate)	1.0%
Sulphur (as sulphate)	7.0%
Boron (as boric acid)	0.8%
Molybdenum (as sodium molybdate)	0.04%
Cobalt (as cobalt sulphate)	0.04%
Maximum Biuret	0.1%

**PRODUCT CONTAINS  
TOTAL SOLIDS**

**45% W/W WATER  
55% W/W**

### GENERAL INFORMATION

CANOPY MASTER 10 is a mixture of 10 essential plant nutrients. It can be used throughout the growing cycle to maintain a healthy canopy and balanced plant growth. It is especially formulated to provide a high analysis mix of important nutrients. CANOPY MASTER 10 is suitable for application to a wide range of crops and can be applied as a soil or foliar treatment.

### MIXING INSTRUCTIONS

- Shake or agitate before use.
- Do not mix with growth hormones unless specifically recommended.
- Compatible with many foliar nutrients and agricultural chemicals.
- Compatible when mixed in the spray tank with Stoller SETT and SETT Enhanced.

**CAUTION: Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.**

**SHAKE WELL BEFORE MIXING**

**STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT**

SII1114

# Stoller's

# CANOPY MASTER 10

"Keeps Leaves Healthy"



**STOLLER AUSTRALIA PTY. LTD.**  
A.C.N. 065 320 747

P O Box 2534  
Regency Park  
South Australia 5942

Phone (08) 82449588  
Fax (08) 82449222  
www.stoller.com.au

**CONTENTS 20 200 800 1000 LITRES**

MADE IN AUSTRALIA

### CAUTION

Avoid contact with eyes and skin. Avoid breathing the spray mist. Wear protective clothing (long pants, long sleeve shirt, gloves hat and boots) during mixing and handling. After use and before eating, drinking or smoking, wash hands, arms, and face thoroughly with soap and water.

### DIRECTIONS FOR USE

CROP	RATE	COMMENTS
Grape Vines	4 litres per hectare	Apply in multiple applications as a foliar. Space treatments at least 2 weeks apart. Can be applied through fertigation at 4-8 litres per hectare.
Vegetables and Horticultural Crops	2 - 4 litres per hectare	Spray as a foliar or as a soil application in a band alongside the planting row.
Tree Crops Citrus and Almonds (avoid foliar application for Stone Fruit)	2 - 4 litres per hectare	Apply in multiple applications as a foliar. Space treatments at least 2 weeks apart. Can be applied through fertigation at 4-8 litres per hectare. Use only as a fertigation treatment for Stone Fruit.
Field Crops	3 - 9 litres per hectare Minimum Dilution 1:25	Apply as required commencing when plants are young. More concentrate can be used on field crops to facilitate lower volume sprays. Product may also be soil applied in a band.
Greenhouse Crops	100 ml/200 litres	Apply frequently at maximum dilution rates for foliar treatment. Can be soil applied at approximately 500ml - 1L per standard house per week.
Turf	200 ml/ 100 sq meters	Repeat applications to maintain healthy growth.

**Avoid application during high daytime temperatures (above 28 degree C)  
Increase dilution rates when spraying in warm conditions**

### CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No: