

# FLOWER POWER



Stoller AUSTRALIA PTY.LTD.

## ANALYSIS

	W/V
ZINC (Zn)	5.0%
BORON (B)	3.8%
COPPER (Cu)	0.1%
MOLYBDENUM (Mo)	0.02%

WITH STOLLER'S PROPRIETARY CO-FACTORS

## GENERAL INFORMATION

Stoller's Flower Power increases the potential for more fruit set and crop yield by optimising the flowering and pollination of any plant. Based on a complex of micronutrients, Flower Power provides key nutrients that improve auxin production and longevity, critical to the plants flowering stage. During extreme temperatures, it continues to improve the pollination strength of flowers.

## MIXING INSTRUCTIONS

- Do not tank mix with growth hormone products, plant growth regulators or foliar fertilisers unless specifically recommended.

**CAUTION:** Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility. Use caution when spraying tank mixes on crops sensitive to fruit marking.

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

SHAKE WELL BEFORE MIXING

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Sellers liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

BMP0717



Stoller

# FLOWER power

STOLLER AUSTRALIA PTY. LTD.

A.C.N. 065 320 747

1 Creswell Rd Phone (08) 8169 0900  
Largs Bay Fax (08) 8169 0909  
South Australia 5016 www.stoller.com.au

CONTENTS | 5 | 20 | 200 | 1000 LITRES

Made in Australia

## CAUTION

Do not contaminate waterways with the chemical or used containers. Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

CROP	RATE PER HECTARE		COMMENTS
Field Crops	2.5-4 L/ha		Apply when first flower appears
Strawberries, Blueberries etc.	2.5-4 L/ha	10-14 days	Begin when first flower appears. Discontinue when last flower is set
Field Vegetables: Tomatoes, Capsicums, Eggfruit	2.5-4 L/ha	7-14 days	Begin when first flower appears. Discontinue use when last fruit has set
Fruit and Nut trees: Fig, Macadamias	2.5 L/ha	10-14 days	Begin when first flower appears. Use higher rates for crops with larger canopies. Discontinue use when last fruit has set
Mangoes, Avocados			Begin at first flower on panicle. Use higher rates for crops with larger canopies. Discontinue use when last fruit has set
Almonds	2.5-4 L/ha	10-14 days	Begin application when pink buds form. Use higher rates for crops with larger canopies. Discontinue when fruit has sized
Grapevines	2.5-4 L/ha	10-14 days	Begin when fruit starts to form. Discontinue when fruit has sized

## Glasshouse Crops

CROP	RATE PER HECTARE		COMMENTS
Tomatoes, Capsicums, Cucumbers	200mL/200L of water	7-10 days	Begin application when first flower appears. Discontinue use when last fruit is set

Batch No.