

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Stoller's N-LARGE™

## Gibberellic Acid Growth Regulant

ACTIVE CONSTITUENT: 100 g/L GIBBERELIC ACID

For Foliar Spray Application to Certain Varieties of Grapes, Citrus and Prunes to Promote Desirable Harvest Effects.

**Important: read the attached booklet before use.**

UN 1170



Flammable Liquid

UN N. 1170	ETHANOL SOLUTION
PG II	HAZCHEM 2 WE
In an emergency Dial 000 Police or Fire Brigade	

**STORAGE AND DISPOSAL:**

Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre (Ph. 131126).

**PRECAUTION:** Flammable. Avoid concentrations higher than those recommended.

**MATERIAL SAFETY DATA SHEET:** Additional information is listed in the Material Safety Data Sheet.

**EMERGENCY INFORMATION:** Contact Stoller Australia, PO Box 2534, Regency Park, SA 5942  
5-7 Pambula Street, Regency Park, SA 5010. Telephone Number: 08 8244 9588.

D.O.M.

 1 Litre

Batch No.

 500 mL

© Registered trademark of Stoller Australia

© Stoller Australia 2004

APVMA Approval No: 58244/500/0104



**STOLLER AUSTRALIA**  
Box 2534  
Regency Park SA 5942  
5-7 Pambula Street  
Regency Park SA 5010

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

**DIRECTIONS FOR USE**

**Restraint:** Do not use with a non-ionic wetter. NOTE: 10 mL product per 100 mL = 10 ppm. Do not apply to plants under pest, nutritional or water stress.

CROP	RATE/100 L	CRITICAL COMMENTS
<b>CITRUS</b>		Apply in minimum volumes of 5000 L/ha to ensure thorough coverage of fruit. Do NOT use where blemish is a problem. For optimum results ensure the spray tank solution is at a pH of 4.0-4.5. See mixing instructions.
Navel and Valencia Oranges	10-20 mL	<b>For reduction in creasing:</b> Prior to applying Stoller's Gibberellic Acid, remove all previous season Valencia fruit. Apply Stoller's Gibberellic Acid to Navel/Valencia oranges when fruit are between 30-50 mm in diameter (generally January – February). Rates lower than 20 ml: <ul style="list-style-type: none"> <li>• Should be used when it is anticipated that fruit will be harvested for early markets</li> <li>• May be used on applications to late navel selections.</li> </ul>
Navel Oranges	10 mL	To delay rind ageing for late marketing (or when fruit is likely to be stored for more than 2 weeks before sale) and reduce rind blemish for longer life, apply when oranges turn from green to silver (colourbreak). When fruit drop is a problem, apply a stop-drop spray. Stoller's Gibberellic Acid is compatible with common stop-drop sprays containing 2,4-D sodium salt. Use 2,4-D at 10 ppm.
Mandarins	10 mL	To delay rind ageing, for late marketing and to reduce rind blemish, apply at three quarters to full colour.
Grapefruit	10 mL	To delay rind ageing, apply the spray when grapefruit turn from green to silver (colourbreak) for grapefruit to be harvested up to mid-November, or apply the spray in mid-June for grapefruit to be harvested in December or January.
Lemons	10 mL	To delay rind ageing, apply Stoller's Gibberellic Acid 4-6 weeks ahead of fruit maturity.

CROP	RATE/100 L	CRITICAL COMMENTS
<b>GRAPES</b>		
Currants – dried fruit	1 mL + 100 ppm Cycocel	<b>To achieve berry thinning:</b> Apply a single combined application (used commonly in SA and NSW) at 100% capfall. Ensure the spray thoroughly covers the bunches.
	100 ppm Cycocel followed by 1 mL Stoller's Gibberellic Acid.	<b>To achieve berry thinning: (Split applications commonly used in Vic.)</b> (i) Apply Cycocel 7 days after bunch droop. (ii) Apply Stoller's Gibberellic Acid at 80-100% capfall.
	200 ppm Cycocel followed by 1 mL Stoller's Gibberellic Acid.	Use the 200 ppm rate of Cycocel on vigorous vines.
	300 ppm Cycocel followed by 1 mL Stoller's Gibberellic Acid.	Use the 300 mL rate of Cycocel on very vigorous Carina vines only. Ensure the spray thoroughly covers the bunches.
Sultanas – dried fruit	10 mL	<b>To achieve berry thinning:</b> Apply Stoller's Gibberellic Acid when bloom or blossom is at 100% capfall stage (full flowering).
Sultanas – fresh fruit		Prune according to vine vigour – avoid more than 8 canes per vine except in special circumstances. Begin thinning late October. Thin bunches so that only the largest bunch per shoot remains. Do not exceed 30 bunches per vine. Bunch trimming should be carried out before fruit set to reduce the incidence of tight bunches. For adequate coverage of table grapes, apply the product in at least 2250 L/ha directed at the bunch area.

CROP
Sultanas fresh fruit (cont.)
Early Madeline
Perlette
Flame seedless

	CROP	RATE/100 L	CRITICAL COMMENTS
used roughly	Sultanas – fresh fruit (cont.)	10 mL	<b>To achieve bunch elongation (stretch):</b> Apply when bunches are half to two-thirds of their final length (when bunches are between 10-15 cm in length). This application is usually applied 10-14 days before the first sign of bloom.
		10 mL	<b>To achieve thinning, two separate applications of 10 mL within the same season are required:</b> Apply the first application of 10 mL at 40% capfall.
		10 mL	Apply the second application of 10 mL at 80% capfall (usually 2-3 days later).
		30 mL	<b>To achieve an increase in berry size, two separate applications of 30 mL within the same season are required:</b> Apply the first application of 30 mL when the smallest berries are 4 mm in diameter, and the largest berries are up to 6 mm dia. Berry shatter may be incomplete at this larger size.
		30 mL	Apply a second application of 30 mL 5 to 7 days later. Trim bunches within two weeks of shatter to leave 3-4 shoulder sprigs. All spray timing stages should be shatter on the top part of the bunch, as the bottom is removed at trimming.
		Early Madeline	20 mL
Ensure	Perlette	12 mL	<b>To achieve berry thinning:</b> Apply at 70% capfall.
		20 mL	<b>To achieve an increase in berry size:</b> Following the 12 mL application for thinning, apply at 20 mL, when berries reach 4-5 mm diameter. Trim bunches as required.
1 bloom	Flame seedless	10 mL	<b>To achieve berry thinning:</b> Apply at 70% capfall.

CROP	RATE/100 L	CRITICAL COMMENTS
Flame seedless (cont.)	30 mL	<b>To achieve increase in berry size, two separate applications of 30 mL within the same season are required:</b> Apply the first application of 30 mL when berries have reached 7-9 mm in diameter.
	30 mL	Apply the second application of 30 mL when the berries have reached 9-10 mm in diameter.
	<b>PRUNES</b>	10 mL

NOT TO BE USED FOR ANY OTHER PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED

### GENERAL INSTRUCTIONS

#### FRUIT QUALITY (GRAPES)

**Bunch Elongation (Stretch):** This product should be sprayed onto bunch stems when bunches are half to two-thirds of their final length (usually 10-15 cm), causes them to grow longer than normal and may prevent over-tightness of bunches. This application is usually applied 10-14 days before the first sign of bloom.

**Thinning:** If sprayed on flowers as they are beginning to open, this product reduces the number of berries on the bunch, i.e., it has a thinning effect.

**Berry size:** This product increases the size of berries when it is applied after the commencement of flowering. The greatest effect occurs when applied at shatter stage.

#### MIXING

Prepare a concentrated solution in 1 to 5 L of water with sufficient product for the required vat volume and strength of spray, add solution to water in the spray vat and agitate.

Citrus only: Ensure the spray mix is in the pH range of 4.0-6.0; however, optimum results occur when the spray mix is in the pH range of 4.0-4.5. Use a portable pH meter or calibrated pH strips to determine the spray mix pH. Sample 2 or 3 times and average the reading. Adjust high pH's with a suitable acidifying solution and recheck the pH after 5 minutes agitation.

**Wetting agent:** Add a maximum of 10 mL per litre of a non-ionic spreader. Citrus only: If using an adjuvant with spreader, do not add additional spreader.

**Precautions**

Do not store prepared solutions for longer than one day.

**CROP MANAGEMENT**

**(Fresh Sultanas):** Prune according to the vigour of the vine. Avoid exceeding 8 canes except in special circumstances. Commence thinning in late October. Thin bunches to leave one bunch per shoot (the largest). DO NOT exceed 30 bunches per vine. Bunch trimming should be carried out after fruit set to reduce the incidence of tight bunches.

**(Citrus):** To maximize product effectiveness and fruit quality, ensure good penetration of spray by skirting and pruning the inside of trees. Hand thinning of fruit may also be of benefit. Even trees with a history of little creasing may require treatment in "heavy crop" years or with age.

Use of Stoller's Gibberellic Acid, particularly at higher rates (20 ppm) can delay colouring by 1-2 weeks, early in the season.

**APPLICATION**

**(Fresh Sultanas):** Make sure vines are watered prior to application of this product. Apply in cool conditions or at night. Where this product is used for dried fruit production, drive-past overall spraying is adequate. For fresh fruit production good results depend on the thorough wetting of bunches and spraying at the correct stage.

**If the bunches are missed they will not react to gibberellic acid.**

FOR RECOMMENDATIONS FOR OTHER GRAPE VARIETIES CONSULT YOUR LOCAL DEPARTMENT OF AGRICULTURE.

**(Citrus):** Spray when the weather is cool, e.g. in the morning or after an irrigation in the afternoon. Avoid application within 4 weeks of any oil spray as the oil restricts GA uptake. Typical water volumes are 5,000 L for small trees, 7,500 L for medium trees and 10,000 L for large trees. For creasing-reduction sprays to be effective, trees must be sprayed to point of runoff.

**COMPATIBILITY:**

Stoller's Gibberellic Acid can be combined in the spray vat with 2,4-D as a cling spray, as well as with products containing Cycocel, certain insecticide, fungicide or nutritional sprays. Always do a small test mix to check compatibility before spraying larger areas. Always check the label instructions for all products used.

**PROTECTION OF CROPS, NON-TARGET PLANTS**

Do NOT apply under adverse weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops.

**PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT:**

Do NOT contaminate streams, rivers or waterways with the chemical or used containers.

**EXCLUSION OF LIABILITY**

Unless otherwise expressly stated in writing neither Stoller Australia Pty Ltd nor the distributor has any knowledge or the particular use to which the buyer proposes to put this product. In purchasing this product the buyer must rely solely upon his own skill and judgment as to its suitability for the particular purpose for which it is required. Except to the extent that exclusion or denial of liability is prohibited under the Trade Practices Act or any relevant state legislation, the Companies and the distributor expressly exclude any warranty as to the quality or fitness of any goods sold for any purpose whatsoever and deny all responsibility in contract tort negligence or otherwise for any harm or damage resulting from the use of such goods or from acting on the advice or recommendations as to such use given in good faith by any representative of the Companies or the distributor. If these conditions are unacceptable to the buyer, the goods should be returned to Stoller Australia Pty Ltd unopened within seven (7) days for refund of purchase price.

© Registered Trademark of Stoller Australia, PO Box 2534, Regency Park, SA 5942. 5-7 Pambula Street, Regency Park, SA 5010

APVMA Approval No. 58244/0104

D.O.M.

Batch No

ledge or  
ly solely  
ot to the  
islation,  
i for any  
damage  
in good  
yer, the  
rice.

SA 5010

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Stoller's N-LARGE™

## Gibberellic Acid Growth Regulant

ACTIVE CONSTITUENT: 100 g/L GIBBERELIC ACID

For Foliar Spray Application to Certain Varieties of Grapes, Citrus and Prunes to Promote Desirable Harvest Effects.

**Important: read the attached booklet before use.**

UN 1170



Flammable Liquid

UN N. 1170	ETHANOL SOLUTION
PG II	HAZCHEM 2 WE
In an emergency Dial 000 Police or Fire Brigade	

#### STORAGE AND DISPOSAL:

Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If no recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SAFETY DIRECTIONS:** Will irritate the eyes and skin. Avoid contact with eyes and skin. When opening the container and preparing the spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre (Ph. 131126).

**PRECAUTION:** Flammable. Avoid concentrations higher than those recommended.

**MATERIAL SAFETY DATA SHEET:** Additional information is listed in the Material Safety Data Sheet.

**EMERGENCY INFORMATION:** Contact Stoller Australia, PO Box 2534, Regency Park, SA 5942  
5-7 Pambula Street, Regency Park, SA 5010. Telephone Number: 08 8244 9588.

D.O.M.

Batch No.

- 1 Litre  
 500 mL

© Registered trademark of Stoller Australia  
© Stoller Australia 2004  
APVMA Approval No: 58244/500/0104



**STOLLER AUSTRALIA**  
Box 2534  
Regency Park SA 5942  
5-7 Pambula Street  
Regency Park SA 5010

# PROOF

## THIS PROOF IS TO BE CHECKED FOR ACCURACY

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors** and **Dieline**.

**Authorized signature** accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.








**PLEASE NOTE:** Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

**THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.**

*Dieline does not print.*

 **Cimarron Label**  
Experts in Extended Text Labeling

4201 North Westport Ave. • Sioux Falls, SD 57107  
Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE	JOB NUMBER	CUSTOMER
10/25/13	71116	STOLLER
LABEL SIZE	LEAFLET FOLDED SIZE	LEAFLET FLAT SIZE
3.875" X 5"	3.875" X 4.25"	3.875" 16.75"
LABEL COLORS	LEAFLET "OUT" COLORS	LEAFLET "IN" COLORS
 348	 300	 185
 348	 300	 185
 348		
PATTERN VARNISH: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

Form: CS 006L - 11/8/2011

**ARTWORK IS APPROVED**

**REVISED PROOF NEEDED**

**WE CANNOT PROCESS  
THIS ORDER WITHOUT AN  
AUTHORIZED SIGNATURE**

Signed \_\_\_\_\_ Date \_\_\_\_\_