

# SETT ENHANCED



Stoller.AUSTRALIA.PTY.LTD.

## ANALYSIS

	W/W
CALCIUM (Ca as chloride)	11.0%
BORON (B)	1.4%

## WITH STOLLER'S PROPRIETARY CO-FACTORS

Derived from: Calcium Chloride, Boron, Polyamines and Complexing Agents.

## PRODUCT CONTAINS

40% W/W WATER

## TOTAL SOLIDS

60% W/W

## GENERAL INFORMATION

SETT ENHANCED is a highly available calcium product including boron and formulated with Stoller's Proprietary Co-Factors. Calcium has a role as a messenger in the plant and boron is known to extend the life of certain key hormones in the plant. The Nutrient - Co-Factor combination in SETT ENHANCED can help with:

- Stronger pollination
- Less impact from cold or hot condition on fertility
- Reducing misshapen fruit in adverse weather
- Better seed/seed trace development
- Stronger flowering in adverse conditions

## MIXING INSTRUCTIONS

- **Do not tank mix with growth hormone products, plant growth regulators or foliar fertilisers unless specifically recommended.**
- SETT ENHANCED can be tank mixed and applied with many commonly used insecticides and fungicides.

**CAUTION: Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility. Use caution when spraying tank mixes on crops sensitive to fruit marking.**

## SHAKE WELL BEFORE MIXING

## STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

BMP0717



Stoller

# SETT ENHANCED

STOLLER AUSTRALIA PTY. LTD.

A.C.N. 065 320 747

1 Creswell Rd  
Largs Bay SA 5016

Phone: 08 8169 0900  
Fax: 08 8169 0909

www.stoller.com.au

MADE IN AUSTRALIA

CONTENTS 20 200 800 1000 LITRES

## CAUTION

Do not contaminate streams, rivers or waterways with the chemical or used containers. Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

## DIRECTIONS FOR USE

It is more effective to spray small amounts with greater frequency. Spray enough water to achieve thorough coverage of the entire plant.

CROP	RATE/HECTARE		COMMENTS
Field Crops	2.5-5 L/ha	7-14 days	Begin application when first flower appears. Discontinue use when last fruit, boll or pods are set
Berries and Strawberries	5-7 L/ha	10-14 days	
Leafy Vegetables	2.5-5 L/ha	7-14 days	Begin application when crop goes into rapid growth' stage, for when tissue starts to develop. Discontinue when rapid growth ceases or when storage tissues finishes bulking
Field Vegetables tomatoes, capsicums, eggfruit	5 L/ha	7-14 days	Begin application when first flower appears. Discontinue when last fruit is set
Tree Fruit and Nuts	5-7 L/ha	10-14 days	Begin when buds first form. Discontinue when fruit has sized
Grapevines	5 L/ha	10-14 days	Begin when fruit starts to form. Discontinue when fruit has sized

## Glasshouse Crops

CROP	RATE		COMMENTS
Tomatoes Capsicums Cucumbers		7-10 days	Begin when first flower appears. Discontinue use when last fruit is set

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Sellers liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No.