

## SP ZINC EDTA Chelate

**ANALYSIS** w/v  
Zinc (Zn) .....7.5% w/v

Derived from Zinc Oxide and EDTA Chelate

### GENERAL INFORMATION

•SP ZINC EDTA Chelate is a foliar or soil applied micronutrient and is non-phytotoxic when used as directed. It is completely available and absorbed by the leaf surface or the root system. SP ZINC EDTA Chelate can be mixed with liquid fertiliser solutions. When mixing SP ZINC EDTA Chelate with liquid fertiliser agitate to thoroughly disperse the trace elements. This can be achieved by adding the product to the mixing batch tank or inducting through a pump. When mixing in batch tank add product to water nitrogen solution before adding phosphate material.

### MIXING INSTRUCTIONS

SP ZINC EDTA Chelate is compatible with many chemicals including MCPA and 2,4-D Amine.

A small amount of wetting agent or surfactant can be added (refer manufacturer's label).

### MIXING SEQUENCE FOR FOLIAR APPLICATION:

1. Water 2. SP ZINC EDTA Chelate 3. Pesticide

**CAUTION: Always Conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.**

**SHAKE WELL BEFORE MIXING**

SI10311

# Stoller's

# SP ZINC EDTA Chelate



**STOLLER AUSTRALIA PTY. LTD.**  
A.C.N. 065 320 747

P O Box 2534 Phone (08) 8440 2420  
Regency Park Fax (08) 8440 2421  
South Australia 5942 www.stoller.com.au

CONTENTS 20 200 800 1000 LITRES

MADE IN AUSTRALIA

### CAUTION

Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist. Use gloves and goggles.

### SP ZINC EDTA Chelate FOLIAR SPRAY

CROP	RATE	COMMENTS
Grain Cereals Wheat, Barley	1 - 3 litres per hectare	Apply between 3 expanded leaf stage through to flowering.
Maize, Corn, Rice, Legumes	1 - 3 litres per hectare	Apply between 3 expanded leaf stage through to flowering.
Canola	1 - 3 litres per hectare	Apply at early emergence, repeat as required.
Cotton	1 - 3 litres per hectare	Apply at early emergence, repeat as required.
Seed Treatment (Wheat)	5 - 7 litres per tonne of seed	Spray using 2 nozzles into auger tube approx. 60 cm apart for thorough coverage. Calibrate to achieve correct application without over-wetting.
Seed Treatment (Smaller Seeds Like Canola)	Up to 20 litres per tonne of seed	
All Other Crops	1 - 3 litres	Commence early. Repeat applications to maintain healthy growth.

*Avoid application in high daytime temperatures (above 28 degree C).  
Increase dilution rates when spraying in warm conditions.*

### CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No: