WL COPPER Chelate

ANALYSIS W/N
COPPER (Cu) 5%
SULPHUR (S) 4%

Derived from: Copper Sulphate and Phenolic Acid Chelate

PRODUCT CONTAINS 82% W/W WATER TOTAL SOLIDS 18% W/W

GENERAL INFORMATION

WL COPPER Chelate is a foliar or soil applied micronutrient and is non-phytotoxic when used as directed. It is completely available and absorbed by the leaf surface or the root system because of the natural complexing properties of its phenolic acid chelating complex

Copper is important for early growth and pollination. Stoller's Chelated Copper can enhance fruit colour and anyhocyanin levels.

MIXING INSTRUCTIONS

WL COPPER Chelate will disperse in water with little agitation. Many pesticides can be added and applied while spraying.

When foliar spraying WL COPPER Chelate through conventional sprayers, use a minimum of 200 litres of water per hectare. When spraying with low volume equipment, 50 litres of water per hectare is usually sufficient. If less water is used, slight burning of the foliage may occur. A maximum of 1.5 litres per hectare per application is recommended. Aerial applications should not exceed 1 litre per 8 litres of water.


WL COPPER Chelate is manufactured to rigid controls at the highest possible concentration for effective usage. Any residue in this pail is water soluble. Rinse the containers with water and add solution to spray tank.

CAUTION: Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING
AVOID SPRAYING WHEN HIGH TEMPERATURES PREVAIL

STORE OUT OF DIRECT SUNLIGHT

SII1215





Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

RECOMMENDATIONS FOR USE

Almonds, apples, avocado, barley, beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, grapes, lettuce, lucerne, melons, milo, nectarines, peaches, peanuts, pears, pecans, peppers, plums, potatoes, prunes, rice, soybeans, sorghum, strawberries, sugar beets, sugar cane, sunflowers, sweet corn, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

DIRECTIONS FOR USE

SEASONAL (TOTAL) RATES FOR SOIL APPLICATION

Maintenance application......2 litres per hectare

Heavy micronutrient deficiency8-10 litres per hectare

FOLIAR SPRAY

FULIAR SPRAY					
CROP	RATE	COMMENTS			
Vegetables	1-1.5 litres per hectare each spray	Incorporate WL COPPER Chelate into the normal spray program.			
Trees, Vines, Nuts	1-1.5 litres per hectare in a dilute spray	If concentrated spray is used, increase Fruit and the concentration in direct proportion to dilution. If tree crops are sprayed several times a year, the amount can be split into more numerous applications. Split applications are considered more beneficial than single applications.			
Field Crops	1-1.5 litres per hectare	Apply recommended rates when plants are young. This is the time when most severe deficiencies occur.			

LIQUID FERTILIZER

Mix WL COPPER Chelate with liquid fertilizer agitating to thoroughly disperse the trace elements. This can be achieved by adding the product to the mixing batch tank or inducting through a pump. When mixing in batch tank, add product to water or nitrogen solution before adding phosphate material.

CONDITIONS OF SAL

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No:		



