

BIO-FORGE®	
ANALYSIS	%W/V
NITROGEN (as total)	2.5%
POTASSIUM (as hydroxide)	3.0%
COBALT	0.25%
MOLYBDENUM	0.35%

CAUTION

Always conduct a “Jar Test” using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT



WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Excessive use of molybdenum (Mo) can be harmful to stock.

Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.



CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller’s control, Seller’s liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



Bio-Forge®

LESS STRESS MORE PROFIT



Stoller Australia
1 Creswell Rd
Largs Bay
South Australia 5016
Phone (08) 8169 0900
Fax (08) 8169 0909
www.stoller.com.au

A.C.N. 065 320 747
MADE IN AUSTRALIA

CONTENTS	1	5	10	LITRES
----------	---	---	----	--------

A Corteva Agriscience Business

BIO-FORGE®

Bio-Forge® regulates specific genes associated with root development and stress ethylene production to allow better nutrient uptake, resulting in improved yields. It is both preventive and curative, allowing growers the flexibility they need to react to plant stress.

DIRECTIONS FOR USE

Apply one application at the start of the season. Reapply if needed.

CROP	RATE
All Crops	1.2 L/ha
Apply immediately before or after a stress event.	
Broadacre Field Crops	125-500 ml/ha
Multiple applications. Refer to your Stoller agronomist for timings.	
Canola, Sorghum, Corn, Cotton, Grain and Legumes	2.5 Litres per tonne of seed
Apply directly to the seed with fungicide.	
Horticulture	300ml-600ml/ha
Multiple applications, to a maximum of 2L/ha per season	
Grape Vines	1.2 L/ha
Apply before or after a stress event.	
Nurseries and Protected Cropping eg. Tomatoes, Capsicums, Eggplant	120ml/100L of water
Apply before or after a stress event.	

BATCH NO: