

## STOLLER'S FOLI-ZYME

ANALYSIS	%W/V
NITROGEN (as total)	15.0%
POTASSIUM (as total)	3.0%
CALCIUM (as total)	4.0%
MAGNESIUM (as total)	1.2%
BORON	0.25%
COBALT	0.01%
COPPER	0.14%
MANGANESE	0.1%
MOLYBDENUM	0.024%
ZINC	0.9%
<i>With Proprietary Co-Factors</i>	
PRODUCT CONTAINS WATER 45% w/w%	
TOTAL SOLIDS 55% w/w%	

**CAUTION:** Always conduct a 'Jar Test' using all products in recommended proportions in water to establish physical compatibility.

**SHAKE WELL BEFORE MIXING**

**STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT**

**WARNING:** Do not swallow. The spray from this product may act as an irritant. Avoid inhalation & contact with the eyes and skin.

Excessive use of molybdenum (mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Mo application may include copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of Mo use with appropriate authorities.

**DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.**



# Foli-Zyme

**ENHANCE HEALTHY GROWTH**



### Stoller Australia

1 Creswell Rd Phone (08) 8169 0900  
Largs Bay Fax (08) 8169 0909  
South Australia 5016 www.stoller.com.au

MADE IN AUSTRALIA

A.C.N. 065 320 747

Contents	20	200	1000	LITRES
----------	----	-----	------	--------

## STOLLER'S FOLI-ZYME

### FOLIAR APPLICATION

For best results spray when crop is actively growing, after irrigation or natural rain fall. Spray early in the morning or late in the afternoon. Midday sprays may not be effective because of excessive moisture loss.

CROP	RATE	COMMENTS
Trees, Fruit, Vines & Nuts	5-9 L/Ha 5-6 times annually beginning with delayed dormant spray. Spray intervals of 14 days are suggested for best results.	Avoid periods when fruit is extremely sensitive to spray damage.
Citrus	5-9 L/Ha every 30 days.	Do not mix with petroleum spray oils or ammoniated copper compounds.
Soil Seeded Field Crops eg. Cotton	5-9 L/Ha when crop is 9cm high.	Continue on 14 day intervals until crop is 14 days from harvest.
Field Row Crops and Vegetables	5-10 L/Ha	To supplement soil applied fertiliser, use a row directed spray. Commence applications when crop has 2-3 leaves or is 20 days old. Continue sprays at 10-14 day intervals.
Lucerne	2.5 - 5 L/Ha	Apply 7-14 days after cutting to enhance growth and dry matter.
Greenhouse	250mls/100L	Apply every 10 to 14 days at first signs of stress.
Turf	500mls/1000 square metres	Repeat every 2-3 weeks until dormancy.

**FERTIGATION** This product can be applied to the soil. Refer to Stoller for rates.

### CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

A Corteva Agriscience Business

Batch No: