

STOLLER'S PHOS MOLY

ANALYSIS	W/V%
NITROGEN (as total)	3%
PHOSPHORUS (as water soluble)	14%
MOLYBDENUM	4%
PRODUCT CONTAINS WATER	46% w/w
TOTAL SOLIDS	54% w/w

CAUTION: Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT



DANGER: May be corrosive to metals. Harmful if swallowed or Inhaled. Causes severe skin burns and eye damage/irritation if contact occurs



WARNING: Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

MIXING INSTRUCTIONS

PHOS MOLY will disperse in water with little agitation. Many pesticides can be added and applied while spraying.

DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.



Phos Moly 3-14-0

FOLIAR NUTRIENT COMPOUND WITH ADDED MOLYBDENUM



Stoller Australia

1 Creswell Rd Phone (08) 8169 0900
Largs Bay Fax (08) 8169 0909
South Australia 5016 www.stoller.com.au
MADE IN AUSTRALIA
A.C.N. 065 320 747

CONTENTS	20	200	1000	LITRES

STOLLER'S PHOS MOLY

RECOMMENDATIONS FOR USE

Almonds, apples, avocado, barley, beans, broccoli, cabbage, carrots, cauliflower, celery, citrus, corn, cotton, grapes, lettuce, lucerne, melons, nectarines, peaches, peanuts, pears, pecans, peppers, plums, potatoes, prunes, rice, soybeans, sorghum, strawberries, sugar beets, sugar cane, sunflowers, sweet corn, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

DIRECTIONS FOR USE

Apply as a foliar spray in a volume of water sufficient for thorough coverage. PHOS MOLY may be applied by conventional ground sprayers, aerial sprayers or overhead irrigation.

RATE	SOIL pH	LEGUMES	NON-LEGUMES	TREES, VINES & VEG
Maintenance Application	5.8 - 6.3	1.0 litres/ hectare	0.5 litres/ hectare	1-2 litres/ hectare
Moderate Deficiency	5.3 - 5.8	2.0 litres/ hectare	1.0 litres/ hectare	3.0 litres/ hectare
Heavy Deficiency	Under 5.3	3.0 litres/ hectare	2.0 litres/ hectare	4.0 litres/ hectare

- Apply PHOS MOLY at 4-5 leaf stage for best results on field crops.
- Apply PHOS MOLY on new foliage for best results on trees, vines and vegetables.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

A Corteva Agriscience Business

BATCH NO: