

STOLLER'S COMO

ANALYSIS	%W/V
COBALT (as total)	1%
MOLYBDENUM	6%
PRODUCT CONTAINS WATER 79% w/w	
TOTAL SOLIDS 21% w/w	

CAUTION: Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING
STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT



WARNING: Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin. Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application.

It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.



CONDITIONS OF SALE
 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



CoMo

6% MOLY

**IMPROVE NODULATION.
 MAXIMISE NITROGEN UTILIZATION**



Stoller Australia
 1 Creswell Rd Phone (08) 8169 0900
 Largs Bay 5016 Fax (08) 8169 0909
 www.stoller.com.au
 A.C.N. 065 320 747

MADE IN AUSTRALIA

CONTENTS	20	200	1000	LITRES
----------	----	-----	------	--------

A Corteva Agriscience Business

STOLLER'S COMO

RECOMMENDATIONS FOR USE

All field crop legumes (soybeans, drybeans, lentils, snap beans, field peas, peanuts) and all forage crop legumes (lucerne, red clover, ladino, clover, lespediza, trifol, subterranean ground peanuts).

CROP	RATE	COMMENTS
Soybeans, cowpeas, dry beans, lentils, snap beans, peanuts & other large seeded legumes:	500 - 800 mls per hectare in 40 litres of water or a volume sufficient for thorough coverage.	Apply at the 4-5 true leaf stage as a foliar spray.
Cereal, Grains & Canola		
Lucerne, red clover, ladino clover, lespediza, trifol & other perennial legumes grown for forage, hay or silage:	500 - 800 mls per hectare in 40 litres of water or a volume sufficient for thorough coverage.	Apply when clover is 4-6 inches tall to assure effective movement into nitrogen fixing nodules. Application should be made no later than 4 weeks prior to cutting.
Horticulture	500 mls per hectare	First application 2-4 weeks from planting. Second application 6-8 weeks from planting.
Blueberries	300 - 500 mls per hectare	Every 1-2 months from late vegetative to harvest stage
Trees & Vines	250 - 500 mls per hectare	Twice per season
CoMo can relieve ethylene stress when applied as a foliar spray at 500ml to 1 litre per hectare		
IN-FURROW		
For ACID SOILS: Apply CoMo at 500ml/ha in-furrow For ALKALINE SOILS: Apply CoMo at 800ml/ha in-furrow		
SEED TREATMENT		
Apply CoMo directly to the seed in the planter box or other suitable container that provides for thorough coating of the seed.		
Soybeans, cowpeas, dry beans, lentils, snap beans, & other large seeded legumes	6 litres per tonne of seed	Apply directly to the seed in the planter box or other suitable container that provides for thorough coating of the seed. If adding inoculant, follow directions on the inoculant label.
Lucerne, clovers & other small seeded legumes	5 litres per tonne of seed	

BATCH NO: