

STOLLER'S CROPPING ZINC

ANALYSIS	%W/V
ZINC (as total)	11.0%
SULPHUR (as total)	5.0%
<i>Together with Phenolic Acid</i>	
PRODUCT CONTAINS 58% w/w WATER	
TOTAL SOLIDS 42% w/w	

MIXING INSTRUCTIONS

- STOLLER'S CROPPING ZINC will disperse in water and can be combined with many agricultural chemicals.
- Check with your local dealer before mixing with seed applied fungicides.

CAUTION: Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT



WARNING: Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.



CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



Cropping Zinc

PHENOLIC CHELATE



Stoller Australia

1 Creswell Rd Phone (08) 8169 0900
Largs Bay Fax (08) 8169 0909
South Australia 5016 www.stoller.com.au

A.C.N. 065 320 747
MADE IN AUSTRALIA

CONTENTS	20	200	1000	LITRES
----------	----	-----	------	--------

STOLLER'S CROPPING ZINC

SOIL APPLICATION (RATE PER SEASON)		
Moderate deficiency	5-7 litres per hectare	
Heavy deficiency	8-12 litres per hectare	
FOLIAR AND SEED TREATMENTS		
CROP	SEED DRESSING	POST EMERGENT
Grain Cereals, Wheat, Barley	3-9 litres per tonne of seed	1-4 litres per hectare each spray. Repeat as required.
Maize, Corn, Rice, Legumes	5-9 litres per tonne of seed	3-7 litres per hectare each spray. Repeat as required.
Canola	8-12 litres per tonne of seed	2-4 litres per hectare each spray. Repeat as required.
Cotton	5-9 litres per tonne of seed	2-3 litres per hectare each spray. Repeat as required.
Field Crops In Furrow	3-15 litres per hectare	Best applied with seed in furrow.
Field Crops General Use Foliar	1-4 litres per hectare	Apply when plants are young. Can be used with Herbicides. DO NOT MIX with AMINES. When using AMINES use Stoller's SP ² Zinc EDTA Chelate.

Best applied when plants are young. Stoller's Cropping Micronutrients are compatible with a wide range of agricultural chemicals. Refer to local agent or Stoller for more information.

Seed application can be applied undiluted or with enough water for even coverage. Product can be used in-furrow, refer to local agent for details.

FOR SEED TREATMENT: SOME ADHERING IN STORAGE MAY OCCUR IF LABEL RATES ARE EXCEEDED OR UNDER ADVERSE CONDITIONS.

A Corteva Agriscience Business

BATCH NO: