

## STOLLER'S MX SPECIAL

ANALYSIS	%W/W
ZINC (as Sulphate)	1.33%
IRON (as Sulphate)	1.33%
BORON (as Total)	0.50%
COPPER (as Sulphate)	0.17%
MANGANESE (as Sulphate)	1.00%
SULPHUR (as Total)	2.50%
MAGNESIUM (as Sulphate)	0.33%
MOLYBDENUM	0.02%

PRODUCT CONTAINS WATER 79.5% w/w  
TOTAL SOLIDS 20.5% w/w

### CAUTION

Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.



### WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

### SHAKE WELL BEFORE MIXING

### STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

DISPOSE OF EMPTY CONTAINER AT AN APPROVED DRUM MUSTER WASTE MANAGEMENT FACILITY.



# MX Special

## FOLIAR NUTRIENT COMPOUND WITH ADDED SECONDARY AND MICRO NUTRIENTS



### Stoller Australia

1 Creswell Rd Phone (08) 8169 0900  
Largs Bay Fax (08) 8169 0909  
South Australia 5016 www.stoller.com.au

A.C.N. 065 320 747

MADE IN AUSTRALIA

CONTENTS	20	200	1000	LITRES
----------	----	-----	------	--------

## STOLLER'S MX SPECIAL

### RECOMMENDATIONS FOR USE

Vegetables, trees, vines, cereals and pasture.

### METHOD OF APPLICATION

#### SOIL APPLICATION

General seasonal application rates when leaf & soil test are not available.

Maintenance application ..... 2 litres per hectare  
Moderate deficiency ..... 2 - 9 litres per hectare  
Heavy micronutrient deficiency ..... 3 - 18 litres per hectare

#### FOLIAR APPLICATION

CROP	RATE	COMMENTS
Vegetables	2-6 litres per hectare each spray	Incorporate MX SPECIAL into the normal spray program.
Trees Fruit, & Nuts	2-6 litres of diluted spray 2-6 litres per hectare	If concentrated spray is used, increase the concentration in direct proportion to dilution. If tree crops are sprayed, several times a year the amount can be split into more numerous applications. Split applications are considered more beneficial than single applications.
Field Crops	2-6 litres per hectare	Apply when plants are young. This is the time when most severe deficiencies occur.

#### LIQUID FERTILISER

Mix MX SPECIAL with liquid fertiliser agitating to thoroughly disperse the trace elements. This can be achieved by adding the product to a fertiliser mix tank or inducting through a pump. When mixing in batch tank, add product to water or nitrogen solution before adding phosphate material.

#### CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

A Corteva Agriscience Business

BATCH NO: