

CANOPY MASTER 10

ANALYSIS	% W/V
NITROGEN (as urea)	5.0%
MAGNESIUM (as sulphate)	3.0%
ZINC (as sulphate)	2.0%
MANGANESE (as sulphate)	2.0%
IRON (as sulphate)	1.0%
COPPER (as sulphate)	1.0%
SULPHUR (as sulphate)	7.0%
BORON (as boric acid)	0.8%
MOLYBDENUM (as sodium molybdate)	0.04%
COBALT (as sulphate)	0.04%
Maximum Biuret	0.1%

Gluconic Acid Chelate

PRODUCT CONTAINS

TOTAL SOLIDS 55% w/w

WATER 45% w/w

WARNING: Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Mo application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of Mo use with appropriate authorities.

CAUTION: Always conduct a 'Jar Test' using all products in recommended proportions in water to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

MIXING INSTRUCTIONS

- Compatible with many foliar nutrients and agricultural chemicals.
- Compatible when mixed in the spray tank with Stoller SETT/CaB & SETT Enhanced.



Stoller®

CANOPY MASTER 10

“Keeps Leaves Healthy”



STOLLER AUSTRALIA PTY. LTD.

A.C.N. 065 320 747

1 Creswell Rd

Phone (08) 8169 0900

Largs Bay

Fax (08) 8169 0909

South Australia 5016

www.stoller.com.au

MADE IN AUSTRALIA

CONTENTS	20	200	1000	LITRES
----------	----	-----	------	--------

WARNING

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation & contact with the eyes and skin.

DIRECTIONS FOR USE

CROP	RATE	COMMENTS
Grapevines	Foliar: 4 L/Ha	Multiple applications in 2 - 4 week intervals.
	Fertigation: 4-8 L/Ha	
Vegetables and Horticultural Crops	2-4 L/Ha	Spray as a foliar or as a soil application in a band alongside the planting row.
Tree Crops Citrus and Almonds	2-4 L/Ha	Multiple applications in 2 - 4 week intervals. Use only as a fertigation treatment for Stone Fruit.
	Fertigation: 4-8 L/Ha	
Field Crops	3-9 litres per hectare Minimum Dilution 1:25	Commence applications during early crop growth. Product may also be soil applied in a band.
Greenhouse Crops	100 ml/200 litres	Apply frequently at maximum dilution rates for foliar treatment. Can be soil applied at approximately 500ml-1L per standard house per week.
Turf	200 ml/ 100 sq meters	Apply multiple applications at 14 day intervals.

Avoid application during high daytime temperatures (above 28°C)

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product.

2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller assumes no liability from handling, storage and use of this product.

BATCH NO: