

# SUGAR MOVER



## ANALYSIS

	W/W
BORON (B)	10.0%
MOLYBDENUM (MO)	0.13%

Derived from: Boric Acid and Sodium Molybdate

Formulated with polyamine complexing agents and Stoller's Co-Factors

**PRODUCT CONTAINS** 35% W/W Water

**TOTAL SOLIDS** 65% W/W

## GENERAL INFORMATION

SUGAR MOVER is a unique foliar nutrient treatment, which will provide boron and molybdenum to a growing plant. It also contains growth co-factors, which help with plant balance. When applied as a foliar, the product will help with the redirection of sugars and carbohydrates away from the growing points and to the fruit and the roots of the plant. The treatment will help with evenness of fruit development and the ability of the plant to withstand stress.

SUGAR MOVER is a convenient effective alternative to soluble boron with the added benefit of molybdenum to promote conversion of nitrate nitrogen into more metabolically functional forms. Boron enhances nitrogen utilisation by improving sugar transport and metabolism, auxin metabolism and seed and fruit development.

Because boron is not mobile within the plant, a continuous supply to all growing points is required from germination to maturity.

**CAUTION:** Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

**STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT**

**SHAKE WELL BEFORE MIXING**

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond sellers control, Sellers liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

BMP 1017



# SUGAR MOVER

**STOLLER AUSTRALIA PTY. LTD.**

A.C.N. 065 320 747

1 Creswell Rd Phone (08) 8169 0900  
Largs Bay Fax (08) 8169 0909  
South Australia 5016 www.stoller.com.au

CONTENTS 5 20 200 1000 LITRES

Made in Australia

## CAUTION

Do not contaminate waterways with the chemical or used containers. Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

## DIRECTIONS FOR USE

CROP	RATE PER HECTARE	COMMENTS
Vegetables (Including potatoes)	1.5-2.5 L/ha	Use 2 applications 7-14 days apart
Tree and Vine	1.5-2 L/ha	Use 2 applications 7-14 days apart
Field Crops (including Canola)	1.5-2 L/ha	Apply at the start of vegetative growth, every 7-14 days until flowering commences
Cotton	1.5-3 L/ha	Apply at first flower bud, for 3 applications 7-14 days apart
Soil Application		Can be applied through the drip or banded between the planting lines. Refer to your Stoller advisor for best rates

**Other Notes:** This product contains boron. Always use in conditions with healthy calcium levels in the plant tissue.

**Compatibility:** Can be included in a foliar program with other nutrients and chemicals. ALWAYS CHECK WITH AN ADVISOR BEFORE PROCEEDING. Aim for a minimum dilution with water of 2%. Higher concentrations can be used on field crops.

**First Aid:** If poisoning occurs contact a doctor or Poisons Information Centre. If swallowed and if more than 15 minutes from a hospital, induce vomiting.

Batch No.