

CALCIUM 5X

LIQUID FERTILISER SOLUTION

GUARANTEED ANALYSIS

CALCIUM
WITH STOLLER'S CO-FACTORS

W/V

5.0%

FOLIAR CALCIUM SPRAY FOR OPTIMISING FRUIT DEVELOPMENT & QUALITY

WARNING

Keep out of reach of children. Avoid contact with skin, eyes or clothing. Wash exposed area with water. Rinse equipment after use.

NOTICE

The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller make any warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.



CALCIUM 5X

STOLLER AUSTRALIA PTY. LTD.
A.C.N. 065 320 747

1 Creswell Rd
Largs Bay
South Australia 5016

Phone (08) 8169 0900
Fax (08) 8169 0909
stoller.com.au

CONTENTS 5 20 200 LITRES

MADE IN AUSTRALIA

FOLIAR APPLICATION RATES

CROP	RATE	COMMENTS
Grapes	5 litres/ha	When fruit is 1-2 mm in size (end of flowering)
	2.5 litres/ha	When fruit is 3-4 mm in size
Trees	5 litres/ha	Full Bloom
	2.5 litres/ha	2 weeks following the first spray
Other Fruits	5 litres/ha	When fruit is 1-2 mm in size
	2.5 litres/ha	When fruit is 3-5 mm in size
Multi Fruiting crops (annuals)	1 litre/ha	Commencing at first fruit formation and then every 10-14 days

NOTES

Calcium 5X is best sprayed by itself or with a Stoller nutrient treatment. The second application to table grapes can coincide with a GA spray. Always maintain a maximum 1 % spray solution and avoid spraying where temperatures exceed 25 degree C. The early spray is the most critical during the beginning of cell division.

BATCH NO: