

LIQUID K

ANALYSIS **W/W**
 POTASSIUM (K) 33%

DERIVED FROM: Potassium Phosphate Carbonate.

PRODUCT CONTAINS **55% w/w WATER**
TOTAL SOLIDS **45% w/w**

GENERAL INFORMATION

- Liquid K is designed to be used as a foliar fertiliser to provide potassium at critical periods of need.
- Potassium is needed to load sugars from the leaf tissue into the phloem tissue so that sugars can exit the leaves and travel to the fruit.
- High levels of potassium should be in the leaves before sugars unload to sizing (bulking) fruit.

MIXING INSTRUCTION

- STOLLER'S Liquid K will disperse in water and can be combined with many agricultural chemicals. A jar test for compatibility is recommended in all cases.

WARNING: Avoid use of Liquid K in multiple product tank mixes, especially with crops sensitive to fruit marking.

CAUTION: Because of its higher PH, care should be used when tank mixing with other fertilisers and chemicals. Always conduct a "jar test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

BMP 0420

Stoller's

WARNING:
 Avoid use of Liquid K
 in multiple product tank mixes,
 especially with crops sensitive
 to fruit marking.



LIQUID K



Stoller

1 Creswell Rd
 Largs Bay
 South Australia 5016
 Phone (08) 8169 0900
 Fax (08) 8169 0909
 stoller.com.au

CONTENTS 20 200 800 1000 LITRES

MADE IN AUSTRALIA

CAUTION:

Do not contaminate streams, rivers or waterways with the chemical or used containers. Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

DIRECTIONS FOR USE

It is more effective to spray small amounts with greater frequency. Spray enough water to achieve thorough coverage of the entire plant.

FOLIAR AND SEED TREATMENTS

CROP	RATE	FREQUENCY	COMMENTS
Broadacre, Cotton & Summer Crops, Wheat & Winter Crops	4-6 litres per hectare	7-10 days	Apply as required to correct deficiencies.
Peanuts, Cucurbits, Tomatoes, Beans and Capsicums	4-6 litres per hectare	7-10 days	Apply during fruit filling & ripening.
Vegetables	4 litres per hectare	7 days	Apply during fruit filling & ripening.
Trees, Fruits and Nuts	4-6 litres per hectare	10 days	Apply during fruit filling & ripening.
Wine Grapes and Table Grapes	4-6 litres per hectare	7-10 days	Use prior to veraison through to 2 weeks before harvest.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

BATCH NO: