

WL IRON CHELATE

ANALYSIS	%w/v
IRON (Fe)	5.0%
SULPHUR (S)	4.0%

Derived from: Ferrous Sulphate and Phenolic Acid Chelate

PRODUCT CONTAINS	76% w/w WATER
TOTAL SOLIDS	24% w/w

GENERAL INFORMATION

WL IRON Chelate is a foliar or soil applied micronutrient and is non-phytotoxic when used as directed. It is completely available and absorbed by the leaf surface or the root system because of the natural complexing properties of its phenolic acid chelating agent.

WL IRON Chelate is an excellent acidification agent for lowering high pH in spray tank water to decrease the hydrolysis of most pesticides

WL IRON Chelate is an effective dispersing agent when applied with other foliar spray chemicals.

MIXING INSTRUCTIONS

WL IRON Chelate will disperse in water with little agitation. Many chemicals can be added and applied while spraying. Refer to pesticide label.

When foliar spraying WL IRON Chelate through conventional sprayers, use a minimum of 200 litres of water per hectare. When spraying with low volume equipment, 50 litres of water per hectare is usually sufficient. If less water is used, slight burning of the foliage may occur. A maximum of 4 litres per hectare per application is recommended. Aerial applications should not exceed 1 litre per 4 litres of water. For tree crops and dilute sprays refer to table opposite. WL IRON Chelate is manufactured to rigid controls at the highest possible concentration for effective usages. Any residue in the pail is water soluble. Rinse with water and add solution to spray tank.

CAUTION: Always conduct a "Jar Test" using all products in proper proportion in order to establish physical compatibility.

SHAKE WELL BEFORE MIXING

STORE IN A COOL PLACE OUT OF DIRECT SUNLIGHT

BMP 0420

Stoller's



IRON Chelate



Stoller

1 Creswell Rd
Largs Bay
South Australia 5016

Phone (08) 8169 0900
Fax (08) 8169 0909
www.stoller.com.au

Contents 10 20 200 800 1000 LITRES

MADE IN AUSTRALIA

CAUTION
Do not contaminate streams rivers or waterways with the chemical or used containers. Harmful if swallowed. Avoid contact with skin, eyes & clothing. Wash thoroughly with soap and water after handling. Avoid breathing spray mist.

DIRECTIONS FOR USE

GENERAL APPLICATION RATES WHEN LEAF AND SOIL TEST ARE NOT AVAILABLE (SOIL APPLICATION)

Maintenance application	2 litres per hectare
Moderate deficiency	4-9 litres per hectare
Heavy nutrient deficiency	9-18 litres per hectare

FOLIAR SPRAY

Crop	Rate	Comments
Vegetables	1-2 litres per hectare each spray	Incorporate WL IRON Chelate into the normal program. Frequent applications may be required depending on severity of deficiency.
Trees, Fruits, and Nuts	2-4 litres per hectare	If tree crops are sprayed several times a year, the amount can be split into more numerous applications. Split applications are more beneficial than single applications.
Field Crops	2-4 litres per hectare	Apply recommended rates when plants are young. This is the time when most severe deficiencies occur.

FERTIGATION RATES

Apply to trees and vines through irrigation at 4-6 litres per hectare per application

LIQUID FERTILIZER

Mix WL IRON Chelate with liquid fertilizer agitating to thoroughly disperse the trace elements. This can be achieved by adding the product to the mixing batch tank or inducting through a pump. When mixing in batch tank, add product to water or nitrogen solution before adding phosphate material.

CONDITIONS OF SALE

1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorised to make any warranty, guarantee or direction concerning this product. 2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Batch No: